ED BY DDC -- NO. 4 09605

USNRDL-TR-651 17 May 1963

CILIATED EPITHELIAL CELLS IN NORMAL MURINE INTRAHEPATIC BILE DUCTS

J. W. Grisham

409 602



U.S. NAVAL RADIOLOGICAL DEFENSE LABORATORY SAN FRANCISCO 24, CALIFORNIA

CELLULAR RADIOBIOLOGY BRANCH G. F. Leong, Head

BIOLOGICAL AND MEDICAL SCIENCES DIVISION E. L. Alpen, Head

ADMINISTRATIVE INFORMATION

This work was accomplished under the Bureau of Medicine and Surgery Task MR005.08-5201, Subtask 1, Technical Objective AW-6, as described in the U. S. Naval Radiological Defense Laboratory Annual Report to the Bureau of Medicine and Surgery (OPNAV FORM 3910-1) of 31 December 1962, and is listed in the U.S. Naval Radiological Defense Laboratory Technical Program Summary for Fiscal Years 1963-1965 of 1 November 1962 under Program A3, Problem 3, entitled "Studies Related to Radiological Casualty Evaluation." This study was supported through funds provided by the Bureau of Medicine and Surgery, and the Defense Atomic Support Agency under NWER Program A4c, Subtask 03.035.

AVAILABILITY OF COPIES

Requests for additional copies by agencies or activities of the Department of Defense, their contractors certified to DDC (formerly ASTIA), and other government agencies or activities should be directed to the Defense Documentation Center for Scientific and Technical Information, Arlington Hall Station, Attn: TIPCR, Arlington 12, Virginia.

Eugene P. Cooper
Eugene P. Cooper
Scientific Director

E.B. Roth, CAPT USN
Commanding Officer and Director

12ND NRDL P1 (3-63)

t. D. Oloto

ABSTRACT

Rare cilia have been noted on cells of normal murine intrahepatic biliary epithelium. In general structure these cilia resemble typical kinocilia but the precise internal structure of the ciliary process has not been ascertained. The function or implication of the presence of cilia on these cells is not absolutely known; they may represent embryological remnants without function.

SUMMARY

The Problem:

Many epithelia are composed of cells containing cilia that function as organs of locomotion to propel materials along the epithelial surface. Studies with the electron microscope have demonstrated occasional cilia on a variety of other types of cells which previously were not suspected to possess these structures. Since their discovery on cells in which they apparently have little locomotive importance, conjecture has arisen as to their function and implication. This report discusses the general problem of function, specifically in relation to cilia on intrahepatic biliary epithelium in normal rats.

The Findings:

Kinocilia, which apparently were typical, were found on epithelial cells of intrahepatic biliary ducts of normal rats. Previous reports have described these structures on abnormally proliferated biliary duct cells and suggestions have been made that the cilia had developed as a "metaplastic change" of the centriole. Since cilia have now been

found on normal biliary epithelial cells, it is apparent that their presence on abnormally proliferated cells does not represent metaplastic change. It is concluded that cilia on bile duct epithelium, as well as on certain other types of cells on which they are rarely found, possibly have no functional importance and may merely represent embryological remnants.

INTRODUCTION

Prior to use of the electron microscope, ciliated cells in bile ducts of mammals had not been described (1). In recent studies using this instrument cilia have been noted on epithelial cells of proliferated intrahepatic bile ducts in rats with various experimental alterations (2, 3, 4) and in human beings with extrahepatic biliary obstruction (3). In these studies ciliated cells were not noted in normal intrahepatic bile ducts in these species although the possibility of their existence was pointed out (3). This report describes the rare occurrence of ciliated cells in small interlobular and intralobular bile ducts of normal adult rats and discusses their possible significance.

MATERIALS AND METHODS

Livers from many untreated normal rats, both male and female, of the Wistar albino strain were examined as controls during the course of many investigations of experimental alterations of hepatic ultrastructure. Small blocks of tissue were fixed in 1% osmium tetroxide in veronal buffer or in Dalton's chromate buffer at pH 7.6. Fixation was carried out at 5°C for 1 hour. Dehydration was accomplished by changes to progressively ascending concentrations of ethyl alcohol solution at intervals of 5 to 10 minutes, and tissues were embedded in Epon 812 by the method of Luft (5). Thin sections were cut on a Porter-Blum ultramicrotome equipped with a glass knife and were examined in an RCA EMU 2B electron microscope after "staining" on the grid with saturated aqueous solution of uranyl acetate.

RESULTS

Cilia were rarely encountered on cells of normal murine intrahepatic biliary epithelium; only six ciliated cells were noted in many
hundreds of sections studied during a period of five years. Only one
cilium was noted on any section of a biliary epithelial cell. In
general structure the cilia resembled typical kinocilia (Figs. 1 and
2), measuring about 250 mu in diameter, having a basal body of approximately 150 by 300 mu and, in most instances, possessing an associated
centriole of similar dimensions. The length and precise internal
structure of the ciliary process could not be ascertained because
favorable longitudinal sections and cross sections were not found in
the few instances in which ciliated cells were encountered. All
cilia seen were located on the lumenal border of the cell and pro-





Fig. 1 A portion of an intralobular bile duct from a normal rat showing a cilium projecting into the lumen of the duct.

Magnification X 13,000.

Fig. 2 An enlargement of the outlined area. The ciliary process (CP) is sectioned tangentially. Basal body (BB) and centriole (C) are shown. Also indicated are numerous microvilli (MV). The lumen of the duct is at the top in both photographs. Magnification X 28,000.

jected into the duct lumen. Cilia were not seen on hepatic parenchymal cells.

DISCUSSION

Studies by electron miscroscopy have recently demonstrated cilia on several diverse types of cells which were previously not suspected to possess these structures. Epithelial cells of normal murine intrahepatic bile ducts may now be added to this heterogenous group which includes renal tumor cells (6) and cells of normal renal epthelium (7), cells in pancreatic islets (8), neurons (9), Schwann cells of unmyelinated nerves (10), fibroblasts (11), smooth muscle cells (12), cells in the adenohypophysis (13), cells in the basal layer of the epidermis and cells of tumors arising from the latter (14), and proliferated intrahepatic biliary epithelial cells in rats (2, 3, 4) and in human beings (3). The structure of the ciliary processes in these instances has been characteristic of kinocilia with 9 peripheral and 2 central filaments (3, 6) or a variation of this arrangement. In the latter instance the observed cilia have lacked central filaments (8, 10, 12, 13), or possessed modified central filaments (11), or have had aberrant numbers of peripheral filaments in addition to lacking central filaments (2). In other instances, as in the present study,

the internal structure of the ciliary process could not be determined (7, 9). Cilia in proliferated biliary epithelium have been described as having the typical 9-2 filament pattern (3) or an unusual 7-0 filament pattern (2). Cilia have been thought to be single on each type of cell mentioned above except in renal tumors (6) and in some instances of biliary ductal proliferation (3) in which several cilia were noted on single cells.

The significance of the presence of cilia on biliary epithelial cells, as well as on the other types of cells previously mentioned, is a matter of speculation. The function of ordinary kinocilia is movement, but the apparent paucity of these structures in the types of cells noted above would seem to indicate that they are of limited importance in this regard. The lack of uniformity in ciliary structure and the heterogeneity of cell type on which they occur would seem to preclude any function common to the group. Various authors have conjectured that they might have such diverse functions as chemoreception (8, 13), creation of turbulence in extracellular fluid (8, 7), or initiation of mitosis (14), or that they might be embryological remnants without function (8, 11). Others have speculated that their presence indicates

"metaplasia of the centriole" (4) or reversion of the cell to a more primitive developmental state (3). The latter two impressions were derived from the belief that cilia were present only on pathologically proliferated cells and not on the normal counterpart. These viewpoints are not justified in the case of proliferated biliary or renal epithelial cells since cilia are present on normal cells (7), as well as on proliferated cells (2, 3, 4, 6).

Biliary epithelial cells of the young of several non-mammalian vertebrate species and of adult petromyzon and certain adult frogs, toads, and reptiles have long been known to possess cilia (1). It is possible that cilia on normal and proliferated murine biliary epithelium and on the other varied types of cells discussed above merely represent structures retained from embryogenic progenitors and that they confer no distinctive attributes on the cells possessing them.

REFERENCES

- 1. Mann, F. C., in Special Cytology, edited by E. V. Cowdry, Paul B. Hoeber, Inc., New York, 1928, 203.
- 2. Steiner, J. W., Carruthers, J. S., Kalifat, S. R., Expt'l and Molecular Pathol., 1962, vl, 162.
- 3. Grisham, J. W., Porta, E. A., Expt'l Cell Research, 1963, In Press.
- 4. Steiner, J. W., Carruthers, J. S., Lab. Invest., 1963, v12, 471.
- 5. Luft, J. H., J. Biochem. Biophys. Cytol., 1961, v9, 409.
- 6. Mannweiler, K., Bernhard, W., J. Ultrastructure Research. 1957, vl, 158.
- 7. Iatta, H., Maunsbach, A. B., Madden, S. C., J. Biochem. Biophys. Cytol., 1961, vll, 248.
- 8. Munger, B. L., Amer. J. Anat., 1958, v103, 275.
- 9. Taxi, J., Compt. Rend. Soc. Biol., 1961, v160, 1860.
- 10. Palay, S. L., Anat. Rec., 1961, v139, 262.
- 11. Grillo, M. A., Palay, S. L., J. Cell Biol., 1963, v16, 430.
- 12. Sorokin, S., ibid, 1962, v15, 363.
- 13. Barnes, B. G., J. Ultrastructure Research, 1961, v5, 453.
- 14. Wilson, R. B., McWhorter, C. A., Lab. Invest., 1963, v12, 242.

Biology and Medicine

INITIAL DISTRIBUTION

Copies

	NAVY
3 1 2 1 1 3 1 1 1 1 1 1 1 1 1	Chief, Bureau of Ships (Code 210L) Chief, Bureau of Ships (Code 320) Chief, Bureau of Medicine and Surgery Chief of Naval Operations (Op-07T) Chief of Naval Research (Code 104) Director, Naval Research Laboratory (Code 2021) Office of Naval Research (Code 422) Office of Naval Research (Code 441) Office of Naval Research FPO, New York Naval Medical Research Institute OiC, Radiation Exposure Evaluation Laboratory Director, Aviation Medical Acceleration Laboratory U. S. Naval Postgraduate School, Monterey Commander, Naval Ordnance Laboratory, Silver Spring Naval Missile Center (Code 5700) CO, Naval Medical Research Unit No. 2 U. S. Naval Hospital, San Diego CO, Naval Medical Field Research Laboratory, Camp Lejeune ARMY
1 1 1 1 1 1 1 1 1 1 1 1	Chief of Research and Development (Atomic Division) Chief of Research and Development (Life Science Division) Peputy Chief of Staff for Military Operations (CBR) Chief of Engineers (ENGMC-DE) Chief of Engineers (ENGCW) CG, Army Materiel Command (AMCRD-DE-NE) U. S. Army Edgewood Arsenal CG, CBR Combat Developments Agency CO, BW Laboratories CO, Fort McClellan, Alabama Commandant, Chemical Corps Schools (Library) CO, Chemical Research and Development Laboratories Commander, Chemical Corps Nuclear Defense Laboratory Hq., Army Environmental Hygiene Agency CG, Aberdeen Proving Ground

1 CO, Army Medical Research Laboratory Army Medical Research and Nutrition Laboratory (MEDEN-AD) 1 CO, Army Medical Service Combat Development Agency 1 Medical Field Service School. Fort Sam Houston (Stimson Lib.) Brooke Army Medical Center (Dept. Prev. Med.) 1 Director, Surgical Research Unit, Fort Sam Houston 1 1 Director, Walter Reed Army Medical Center 1 Hq., Army Nuclear Medicine Research Detach., Europe 1 CG, Combat Developments Command (CDCMR-V) CG, Quartermaster Res. and Eng. Command 1 Hq., Dugway Proving Ground 3 The Surgeon General (MEDNE) 1 Office of the Surgeon General (Combat Dev.) 1 CO, Engineer Res. and Dev. Laboratory 1 Director, USACDC Nuclear Group 1 CG, Munitions Command CG, Frankford Arsenal CG, Army Missile Command AIR FORCE Assistant Chief of Staff, Intelligence (AFCIN-3B) 1 6 CG, Aeronautical Systems Division (ASAPRD-NS) CO, Radiological Health Laboratory Division 1 Director, USAF Project RAND Commandant, School of Aerospace Medicine, Brooks AFB 1 CO. School of Aviation Medicine, Gunter AFB Air Force Special Weapons Center - SWRBB 1 1 Radiobiological Laboratory Office of the Surgeon (SUP3.1), Strategic Air Command 1 Office of the Surgeon General 1 Commander, Special Weapons Center, Kirtland AFB 1 1 Director, Air University Library, Maxwell AFB Commander, Technical Training Wing, 3415th TTG 2 1 Hq., Second Air Force, Barksdale AFB Commander, Electronic Systems Division (CRZT) OTHER DOD ACTIVITIES 3 Chief, Defense Atomic Support Agency (Library) Commander, FC/DASA, Sandia Base (FCDV) 1 Commander, FC/DASA, Sandia Base (FCTG5, Library) 1 1 Commander, FC/DASA, Sandia Base (FCWT) 2 Office of Civil Defense, Washington Civil Defense Unit, Army Library 2

Armed Forces Institute of Pathology

- 20 Commander, Defense Documentation Center
- 1 Director, Armed Forces Radiobiology Research Institute

AEC ACTIVITIES AND OTHERS

- 1 Research Analysis Corporation
- 1 Director, Division of Biology and Medicine
- 1 NASA, Ames Research Center, Moffet Field
- 1 Aerojet General, Azusa
- 5 Argonne Cancer Research Hospital
- 10 Argonne National Laboratory
- 2 Atomic Bomb Casualty Commission
- 1 AEC Scientific Representative, France
- 1 AEC Scientific Representative, Japan
- 3 Atomic Energy Commission, Washington
- 2 Atomic Energy of Canada, Limited
- 3 Atomics International
- 2 Battelle Memorial Institute
- 1 Borden Chemical Company
- 3 Brookhaven National Laboratory
- 1 Chicago Patent Group
- 1 Colorado State University
- 1 Columbia University (Rossi)
- 1 Committee on the Effects of Atomic Radiation
- 3 Defence Research Member
- 2 duPont Company, Aiken
- 1 duPont Company, Wilmington
- 1 Edgerton, Germeshausen and Grier, Inc., Goleta
- 1 Edgerton, Germeshausen and Grier, Inc., Las Vegas
- 2 General Dynamics, Fort Worth
- 2 General Electric Company, Cincinnati
- 8 General Electric Company, Richland
- 1 General Electric Company, St. Petersburg
- 1 General Scientific Corporation
- 1 Hughes Aircraft Company, Culver City
- l Iowa State University
- l Journal of Nuclear Medicine
- 1 Knolls Atomic Power Laboratory
- 2 Los Alamos Scientific Laboratory (Library)
- 1 Lovelace Foundation
- 1 Martin-Marietta Corporation
- 1 Massachusetts Institute of Technology
- 1 Mound Laboratory
- 2 NASA, Scientific and Technical Information Facility
- 1 National Academy of Sciences
- 1 National Bureau of Standards (Taylor)

```
1.
        National Cancer Institute
1
        National Lead Company of Ohio
1
        National Library of Medicine
1
        New Jersey State Department of Health
1
        New York Operations Office
1
        New York University (Eisenbud)
1
        Office of Assistant General Counsel for Patents
2
        Phillips Petroleum Company
4
        Pratt and Whitney Aircraft Division
2
        Public Health Service, Washington
1
        Public Health Service, Las Vegas
1
        Public Health Service, Montgomery
1
        Reynolds Electrical and Engineering Company, Inc.
1
        Sandia Corporation, Albuquerque
1
        Union Carbide Nuclear Company (ORGDP)
5
        Union Carbide Nuclear Company (ORNL)
1
        Union Carbide Nuclear Company (Paducah Plant)
1
        United Nuclear Corporation (NDA)
1
        U. S. Geological Survey, Denver
1
        U. S. Geological Survey, Menlo Park
1
        U. S. Geological Survey, Naval Gun Factory
1
        U. S. Geological Survey, Washington
1
        U. S. Weather Bureau, Washington
1
        University of California, Davis
3
        University of California Lawrence Radiation Lab., Berkeley
2
        University of California Lawrence Radiation Lab., Livermore
1
        University of California, Los Angeles
1
        University of California, San Francisco
1
        University of Chicago Radiation Laboratory
1
        University of Hawaii
1
        University of Puerto Rico
1
        University of Rochester (Atomic Energy Project)
1
        University of Tennessee (UTA)
1
        University of Utah
1
        University of Washington (Donaldson)
1
        Wayne State University
1.
        Westinghouse Electric Corporation (NASA)
        Westinghouse Electric Corporation (Rahilly)
1
1
        Western Reserve University (Friedell)
25
        Technical Information Extension, Oak Ridge
```

USNRDL

41 USNRDL, Technical Information Division

DISTRIBUTION DATE: 5 July 1963

1	Naval Radiological Defense Laboratory USNRDL-TR-651 CILIATED EPITHELIAL CELLS IN NORMAL MURINE INTRAHEPATIC BILE DUCTS by J. W. Grisham 17 May 1963 16 p. illus. 14 refs. Grisham 17 May 1963 16 p. illus. 14 refs. Rare cilia have been noted on cells of normal murine intrahepatic biliary epithelium. In general structure these cilia resemble typical kinocilia but the ciliary process has not been UNCLASSIFIED	r implication of the presence i; they may represent embryol
	1. Cilia and ciliary motion. 2. Biliary system. 3. Epithelium. 4. Cells (Biology). 5. Rats. 1. Grisham, J. W. 11. Title. 11. MR005.08-5201	sence of cilia on these nbryological remnants
4	Naval Radiological Defense Laboratory USNRDL-TR-651 CILIATED EPITHELIAL CELLS IN NORMAL MURINE INTRAHEPATIC BILE DUCTS by J. W. Grisham 17 May 1963 16 p. illus. 14 refs. UNCLASSIFIED Rare cilia have been noted on cells of normal murine intrahepatic biliary epithelium. In general structure these cilia resemble typical kinocilia but the precise internal structure of the ciliary process has not been	ascertained. The function or implication of the presence of cilia on these cells is not absolutely known; they may represent embryological remnants without function. UNCLASSIFED